

Repeated downsweep vocalizations of the Araguaian river dolphin, *Inia araguaiensis*

Gabriel Melo-Santos,^{1,a)} Sam F. Walmsley,¹ Miriam Marmontel,² Marcelo Oliveira-da-Costa,³ and Vincent M. Janik¹

¹Sea Mammal Research Unit, Scottish Oceans Institute, School of Biology, University of St Andrews, East Sands, KY16 8LB, St Andrews, Scotland, United Kingdom

²Research Group on Amazonian Aquatic Mammals, Mamirauá Sustainable Development Institute, Estrada do Bexiga No. 2584, Tefé, Brazil

³WWF Brazil (World Wide Fund for Nature), Brasília, Federal District, Brazil

ABSTRACT:

Araguaian botos (*Inia araguaiaensis*) are known to produce pulsed as well as tonal sounds. This study documents the first evidence for repetitive sequences of downsweep whistles in botos that appear to be shared between individuals, and the context of their occurrence is investigated. Boat surveys were conducted along the Tocantins River located in the Eastern Amazon over a period of 42 days between 2012 and 2018. Eighty-two groups of Araguaian botos were observed, and 43 h of sound recordings were acquired. 632 downsweep whistles were recorded in 10 encounters. Four of these encounters contained downsweep bouts (21 bouts with ≥ 2 whistles) with short inter-call intervals (bout criterion 50 s) and up to 161 whistles. A statistical relationship was not found between downsweep occurrence and any of the contextual parameters that were investigated, including socializing, travelling, feeding, group size, presence of calves, and socio-sexual displays. The rarity of these signals makes them unlikely candidates for individual or group identification. It is more likely that they are associated with very specific contexts, such as nursing or mating, both of which were rarely observed in this study. Further studies are required to investigate context specificity and elucidate the function of these signals. © 2020 Acoustical Society of America.

<https://doi.org/10.1121/10.0000624>

(Received 12 August 2019; revised 6 January 2020; accepted 9 January 2020; published online 5 February 2020)

[Editor: Brian Branstetter]

Pages: 748–756

I. INTRODUCTION

All toothed whales (Odontoceti) produce pulsed calls for communication, with some species using these as their main social signals, e.g., orcas (*Orcinus orca*; Ford, 1989; Deecke *et al.*, 2010), northern right whale dolphins (*Lissodelphis borealis*; Rankin *et al.*, 2007), narwhals (*Monodon monoceros*; Marcoux *et al.*, 2012), and pilot whales (*Globicephala* species; Sayigh *et al.*, 2013; Pérez *et al.*, 2017). Pulsed calls are discrete signals composed of a series of individual clicks (Schevill and Watkins, 1966). Such pulsed calls often encode social group identity (Deecke *et al.*, 2010; Ford, 1989) but are also used in immediate social interactions such as conflict negotiation (Overstrom, 1983). Many marine dolphins (family Delphinidae) also use whistles in their social communication (Janik, 2005; May-Collado *et al.*, 2007). Whistles are defined as narrowband frequency modulated tonal sounds (Au *et al.*, 2000; Richardson *et al.*, 1995; Tyack, 2000). For most delphinids these sounds help to maintain cohesion of social groups (Janik and Slater, 1998; Janik, 2005), and for some species, such as the bottlenose dolphins (*Tursiops* species), whistles are also used to broadcast individual identity (Janik *et al.*, 2006; Sayigh *et al.*, 1999).

River dolphins are a polyphyletic group that is particularly interesting when studying acoustic communication in cetaceans because riverine habitats are very different from marine ones and may have led to changes in communication strategies. However, comparatively few studies exist on their vocalizations. Previous studies on the susu (*Platanista gangetica*) and the baiji (*Lipotes vexillifer*) were mainly conducted in captivity and therefore focused on very few individuals (Mizue *et al.*, 1971; Wang *et al.*, 1995; Wang *et al.*, 1999; Wang *et al.*, 2006; Xiao and Jing, 1989). Cremer *et al.* (2017) recently described the whistles and burst pulses of wild franciscana dolphins (*Pontoporia blainvillei*) and suggested that they might be used in mother-calf communication. The most commonly studied river dolphin is the boto (genus *Inia*; Amorim *et al.*, 2016; Caldwell *et al.*, 1966; Diazgranados and Trujillo, 2002; May-Collado and Wartzok, 2007; Ding *et al.*, 1995; Ding *et al.*, 2001; Kamminga *et al.*, 1993; Podos *et al.*, 2002; Penner and Murchison, 1970). Initially they were thought to lack whistles in their repertoire (Podos *et al.*, 2002), but later studies showed that whistles do occur, just not as often as pulsed sounds (May-Collado and Wartzok, 2007; Melo-Santos *et al.*, 2019). The discrepancy in these results may be partly explained by geographic and genetic differences. For example, a recent study has discovered that botos in the Tocantins-Araguaia River Basin belong to a different

^{a)}Also at: BioMA—Biology and Conservation of Amazonian Aquatic Mammals, Belém, Brazil. Electronic mail: gmad@st-andrews.ac.uk

species, the Araguaian boto (*Inia araguaiaensis*), than those found elsewhere (Hrbek *et al.*, 2014).

Melo-Santos *et al.* (2019) described a variety of sounds produced by this new species, including whistles and pulsed calls. Araguaian boto calves were also found to produce bouts of short pulsed calls that were hypothesized to be contact calls for mother-calf communication (Melo-Santos *et al.*, 2019). Repetition introduces redundancy into a signal and thereby increases the probability of transmitting information successfully to a receiver.

Various animal groups produce repeated call sequences, including birds (Catchpole and Slater, 2008), insects (Hedrick, 1986), frogs (Fellers, 1979), and terrestrial mammals (McComb, 1991). In cetaceans, rhythmically repeated calls have been reported for a wide range of species, such as bottlenose dolphins (Jensen *et al.*, 2012; Janik *et al.*, 2013), short and long-finned pilot whales (*Globicephala macrohynchus* and *Globicephala melas*, respectively; Sayigh *et al.*, 2013; Zwamborn and Whitehead, 2017), northern right whale dolphins (Rankin *et al.*, 2007), melon-headed whales (*Peponocephala electra*; Kaplan *et al.*, 2014), Guiana dolphins (Duarte de Figueiredo and Simão, 2009), humpback whales (*Megaptera novaeangliae*; Payne and McVay, 1971), sperm whales (*Physeter macrocephalus*; Watkins and Schevill, 1977), and short-beaked common dolphins (*Delphinus delphis*; Fearey *et al.*, 2019). Repetitive signals might work as mating calls (McComb, 1991;

Smith *et al.*, 2008), individual or group identifiers (Janik *et al.*, 2013; Gero *et al.*, 2016; Zwamborn and Whitehead, 2017), mother-offspring contact (Smolker *et al.*, 1993), territorial defense signals (Fellers, 1979), or even food calls (Janik, 2000). Here, we present the first evidence for repetitive sequences of downsweep whistles in botos that appear to be shared between individuals and investigate the context of their occurrence.

II. METHODS

A. Study animals and area

Dolphins of the genus *Inia* are found in the Amazon, Orinoco, and Tocantins river basins in South America (Best and Da Silva, 1989, 1993; Hrbek *et al.*, 2014; Santos *et al.*, 2012; Santos *et al.*, 2014). These dolphins prefer slow flowing, sheltered waters with large concentrations of prey (Gomez-Salazar *et al.*, 2012b; Martin *et al.*, 2004; Pavanato *et al.*, 2016). They are usually solitary or in mother-and-calf pairs. Larger aggregations are formed for mating and to feed on large schools of fish (Best and Da Silva, 1989, 1993; Gomez-Salazar *et al.*, 2012a; Martin *et al.*, 2008).

Our study was conducted along the Tocantins River located in the Eastern portion of the Amazon (Fig. 1). This river is characterized by clear waters, sandbanks, herbaceous and floating vegetation, and aquatic macrophytes in areas with light penetration (Junk *et al.*, 2011). The greatest




FIG. 1. (Color online) Location of study area and sound recordings of Araguaian botos, *I. araguaiaensis*, in the Tocantins River.

rainfall occurs from November to April, the lowest waters are in September, and the highest waters are in March (Ribeiro *et al.*, 1995). The lower Tocantins River comprises the area between the mouth of the river and the city of Tucuruí, where there is deposition of sediments and floodplains (Goulding *et al.*, 2003; Ribeiro *et al.*, 1995; Santos and Jégu, 1989). The area upstream of the Tucuruí dam is known as middle Tocantins River (Goulding *et al.*, 2003; Ribeiro *et al.*, 1995; Santos and Jégu, 1989). The Tocantins River is heavily impacted by human activities in its vicinity such as the operations of large cities and farms along the banks, fishing, the use of water for irrigation and electrical power generation, the operation of vessels for transport and fishing, and the refinement of ore (Goulding *et al.*, 2003; Ribeiro *et al.*, 1995).

B. Data collection

We conducted 42 days of boat surveys from June 2012 to January 2018. Surveys were conducted around the Capim Island (01°33'48.43"S 48°50'37.81"W) in the mouth of the Tocantins River and between the cities of Baião (02°38'57.26"S 49°40'44.94"W) and Marabá (05°19'21.02"S 49°7'29.02"W), including the reservoir of the Tucuruí Hydroelectric Plant (Fig. 1, Table I). Around the Capim Island we used a 7 m wooden boat with a two-stroke center engine, and for the rest of the surveys we used a 10 m aluminum boat with 40 hp four-stroke outboard engine. During surveys we maintained a boat speed between 5.4 and 8.1 kn in an area up to 200 m from shore. *Inia* dolphins prefer these shallow areas most likely due to increased prey density (Gomez-Salazar *et al.*, 2012a; Pavanato *et al.*, 2016). When possible, we crossed the river once every 10 km to include animals in the middle of the river channel in our recordings.

When a group of dolphins was sighted, we reduced speed, shut the engine off, and began recording when we were approximately 20–50 m from the animals. Recording sessions lasted between 2.53 min and 2.3 h; the recording time depended on our ability to track the animals. If a group avoided the boat approach three times we proceeded searching for another group. In 2012 and 2013 we used an Aquarian Hydrophone System (Anacortes, WA, USA) connected to a Tascam DR-1 recorder (Santa Fe Springs, CA, USA), with a sampling rate of 44 kHz. In 2017 and 2018 we recorded dolphins using a High Frequency Soundtrap (Oceans Instruments, Warkworth, Auckland, New Zealand), with a sampling rate of 576 kHz. During recordings we took notes on group size and age composition, behavior (Table II)

habitat, geographical position, and river state (0–3 in an increasing scale of turbulence). Behavioral sampling of aerial behavior followed the continuous *all event* method (Altmann, 1974). A group was defined as animals in a radius of 100 m from each other; we monitored continuously for group size and changes in composition and took note of the highest number of animals for each encounter. We considered calves as those animals with less than half the size of an adult, also recognizable by their stunted rostrum. Habitat types were classified according to Gomez-Salazar *et al.* (2012a) and Pavanato *et al.* (2016) as follows: main river channel, channel, island, main river margin, lake, streams, and rocky formations. Behavioral states were categorized as (1) travelling, unidirectional movement in any speed; (2) socializing, frequent body contact between animals and surface displays (e.g., leaps, object-carrying), and (3) feeding, repeated diving in the same area, surfacing in multiple directions, and diving with an arched dorsum. As botos are normally slow swimmers (Best and Da Silva 1989, 1993) and the animals we observed were always in motion, thus we did not include a resting category.

C. Acoustic and statistical analyses

We defined downsweeps as tonal signals with the initial frequency higher than the final frequency and no inflection points. An inflection point was defined as a point where the slope of the whistle contour reverses direction (Au *et al.*, 2000). Acoustic analysis was carried out in the software Luscinia.¹ We identified downsweeps with high signal-to-noise ratio, traced their contours, and extracted the following parameters: fundamental frequency, peak frequency range, duration, Wiener entropy, and harmonicity. The last two parameters were added as measurements of noisiness because sounds produced by *Inia* dolphins have, sometimes, noisy components. Then, we used a dynamic-time warping algorithm to compare the measured parameters between calls. This technique allows for extension and compression of a signal along the time axis, ensuring maximum overlapping of the frequency domain when comparing two signals (Buck and Tyack, 1993; Deecke and Janik, 2006; Sakoe and Chiba, 1978). The resulting dissimilarity matrix from the time-warping analysis was converted to Euclidean distances for further statistical analysis. We then ran a nonmetric multidimensional scaling analysis (NMDS). The dimensions of the NMDS were used in a principal component analysis (PCA) for further data reduction. The results were plotted to visualize differences in downsweeps produced by different groups and populations. Inter-call intervals and downsweep parameters were measured in Raven Pro 1.5 (Cornell Laboratory of Ornithology, New York, NY, USA). Downsweep whistle bouts with an inter-whistle interval shorter than 0.05 s were excluded from the analysis, as it appeared that they were produced by more than one individual as indicated by occasional overlap of whistles in such bouts (112 downsweep whistles in 5 bouts excluded).

TABLE I. Summary of surveying effort along the Tocantins River.

Location	Month	Year	Number of surveying days
Capim Island	June	2012	1
	July	2013	3
Baião-Marabá	September	2017	20
	January	2018	18

To investigate the behavioral context of *Inia* downsweeps, we modelled the presence and absence of downsweeps in each recording session using a generalized linear model (GLM; logit link, binomial family)

$$\bar{D} = \frac{\exp(\beta_0 + \beta_1\phi_s + \beta_2\phi_t + \beta_3c + \beta_4g + \beta_5r)}{1 + \exp(\beta_0 + \beta_1\phi_s + \beta_2\phi_t + \beta_3c + \beta_4g + \beta_5r)},$$

$$D \sim \text{Binomial}(\bar{D}),$$

where D represents the probability of one or more downsweeps being produced in a recording session, and \bar{D} represents expected probabilities. Parameters β_1 and β_2 represent the difference in model intercept (β_0) when dolphins were social-

izing (ϕ_s) or travelling (ϕ_t), relative to feeding. The effects of the number of calves present (c) and group size (g) are represented by β_3 and β_4 , respectively. Because our ability to record dolphin sounds varied across a behavioral state, some states were associated with longer recording durations than others (e.g., socializing mean = 65.63 min \pm 8.06, and feeding mean = 28.46 min \pm 3.84). Expecting that a call type should be more likely to be detected with longer recording samples, we simultaneously estimated β_5 , the effect of recording duration (r , mins) to avoid confounding the effects of behavior on downsweep production.

Next, to investigate possible relationships between downsweep production and discrete behavioral events, we fit the following GLM (logit link, binomial family):

$$\bar{D} = \frac{\exp(\varepsilon_0 + \varepsilon_1\phi_b + \varepsilon_2\phi_f + \varepsilon_3\phi_l + \varepsilon_4\phi_o + \varepsilon_5\phi_p + \varepsilon_6\phi_s + \varepsilon_7\phi_t + \varepsilon_8r)}{1 + \exp(\varepsilon_0 + \varepsilon_1\phi_b + \varepsilon_2\phi_f + \varepsilon_3\phi_l + \varepsilon_4\phi_o + \varepsilon_5\phi_p + \varepsilon_6\phi_s + \varepsilon_7\phi_t + \varepsilon_8r)},$$

$$D \sim \text{Binomial}(\bar{D}),$$

where parameters $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_7$ represent the difference in expected probability of a downsweep being produced in a given recording session (\bar{D}) when “body contact,” “flippers, fluke, and belly exhibitions,” “leaps,” “object-carrying,” “penis exhibitions,” “synchronized leaps,” or “tail slaps” occurred, respectively, and ε_8 represents the effect of recording duration (r , mins; see Table II for descriptions of discrete behavior patterns). Both of the above models were also run as a quasi-binomial model to test for overdispersion. All statistical analyses were conducted using R 3.4.2 (R Core Team, 2017).

In order to define downsweep bouts, we used a log survivorship analysis as described by Slater and Lester (1982). We present inter-whistle intervals within bouts, as well as between bouts, and single downsweep emissions.

III. RESULTS

We encountered 82 groups of Araguaian botos, comprising sightings of 385 individuals. Group size varied from 1 to

20 animals (mean = 5 \pm 0.5). Calves were present in 28 groups. We analyzed 43 h of sound recordings from these sightings, identifying 632 downsweeps in recordings from 10 groups; calves were observed in 7 of these groups. The minimum number of downsweeps in an encounter was one, and the maximum number of downsweeps in an encounter was 287 (Table III, Fig. 2). Downsweeps were often produced in bouts with short inter-call intervals. We used 50 s as a bout inter-whistle interval criterion determined by log analysis of the distribution of inter-whistle intervals (Slater and Lester, 1982). We found 21 bouts with the shortest bout consisting of only 2 whistles and the longest of 161 whistles (median = 9 whistles, 25% quantile = 3, 75% quantile = 25). Inter-whistle intervals within bouts ranged from 0.052 to 45.82 s (median = 1.01, 25% quantile = 0.71, 75% quantile = 2.37). Intervals between bouts or single downsweep emissions varied from 51.12 to 1901.65 s (median = 164.89, 25% quantile = 75.51, 75% quantile = 317.20; Fig. 3, Table III). Among the 82 groups we observed in our study, 3 displayed social-sexual behavior (calves present in all these observations), which included

TABLE II. Definition of discrete behavioral events.

Behavioral event	Definition
Body contact	Animals touching each other's bodies
Flippers, fluke, and belly exhibitions	Dolphins swimming sideways or belly-up so that the flippers, belly, and/or fluke are shown above the water surface
Leap	Single animal jumps out of the water exhibiting most of its body
Synchronized leap	Two or more dolphins jump out of the water exhibiting most of their bodies
Object-carrying	Dolphin holding an object (e.g., pebble, vegetation) with its rostrum out of the water
Penis exhibition	Animal swimming belly-up and exhibiting its penis out of water
Tail slap	Dolphin slapping its tail fluke on the water surface one or multiple times

TABLE III. Characterization of all downsweep whistles. The “L” in group ID indicates animals recorded in the Lower Tocantins River, while the “M” stands for recordings of the Middle Tocantins River. Encounters where we detected downsweep bouts have their group ID in bold. Mean and standard error (SE) are presented for acoustic parameters of downsweeps.

Group ID	Behavioral state	Group size	Number of calves	Recording time (min)	Number of downsweeps	Duration (sec ± SE)	Minimum frequency (kHz ± SE)	Maximum frequency (kHz ± SE)	Frequency range (kHz ± SE)	Peak frequency (kHz ± SE)
L1	Feeding	5	1	33.48	161	0.06 ± 0.001	5.661 ± 0.069	16.98 ± 0.13	11.32 ± 0.13	7.04 ± 0.18
L2	Traveling	2	0	4.37	1	0.04	7.27	11.24	3.97	8.58
L3	Socializing	6	1	52.15	44	0.04 ± 0.005	3.52 ± 0.15	5.45 ± 0.16	1.93 ± 0.12	3.97 ± 0.16
M1	Socializing	18	1	138.5	131	0.042 ± 0.002	3.83 ± 0.08	6.09 ± 0.16	2.64 ± 0.12	4.63 ± 0.09
M2	Feeding	8	2	118.15	4	0.03 ± 0.002	3.53 ± 0.51	5.25 ± 1.01	1.72 ± 0.63	3.90 ± 0.44
M3	Socializing	15	1	87.36	287	0.07 ± 0.001	9.19 ± 0.04	13.67 ± 0.05	4.48 ± 0.06	9.95 ± 0.05
M4	Feeding	2	0	18.02	1	0.07	2.66	4.80	2.13	3.37
M5	Socializing	3	0	56.26	1	0.09	8.02	20.42	12.40	8.86
M6	Feeding	8	1	56.85	1	0.02	11.65	17.13	5.48	11.95
M7	Feeding	4	1	125.9	1	0.05	9.20	14.05	4.84	13.08
Total		71	8	691.04	632	0.06 ± 0.001	6.73 ± 0.10	12.32 ± 0.17	5.58 ± 0.14	7.64 ± 0.10

object-carrying, exhibition of the penis, and close body contact between individuals. Of these three groups, two emitted downsweep bouts.

Table III summarises downsweep whistle parameters. In the PCA, the first principal component explained 68.2% of the variance in the downsweep parameters, and the second principal component explained 14.9% of the variance. While there was no clear differentiation between populations (lower and middle Tocantins River), two groups, one from each population, produced very characteristic signals and formed tight clusters on the ordination analysis (Fig. 4).

To investigate the context of downsweep whistling, we conducted GLM analyses. We were unable to confidently identify a behavioral state for 3 of 82 recordings, as the animals were only observed on the surface once. These

recordings were excluded from the GLMs. There was no evidence of overdispersion in either the quasi-binomial behavioral state GLM (overdispersion parameter = 0.99) or the discrete-event GLM (overdispersion parameter = 1.32). Thus, we interpreted parameter estimates from the models of the binomial family. Relative to feeding (the background behavioral state in our model), downsweeps were estimated to be more likely to occur when animals were travelling or socializing, although these estimates were associated with large standard errors (SEs) and were not statistically significant (Table IV). The presence of calves was also included in the analysis but, again, no significant relationship with downsweep production was discovered. Downsweeps were significantly more likely to be detected in long recordings, suggesting that sampling duration was more influential than




FIG. 2. Examples of downsweeps produced by Araguaian botos in the Tocantins River. Labels in the spectrograms indicate the group of botos from which the sound recording is represented. The “L” in group identification (ID) indicates animals recorded in the Lower Tocantins River, while the “M” stands for recordings of the Middle Tocantins River.




FIG. 3. (Color online) Distribution of downsweeps within each recording session. \times 's represent downsweep emissions, and the continuous lines represent the recording time for each group of Araguaian botos.

the behavioral and group measures we included. Similarly, we found that none of the discrete behavioral events that we analyzed had a significant effect on the occurrence of downsweeps (Table V).

IV. DISCUSSION

Downsweep whistles of Araguaian botos were often produced in long bouts with short inter-call intervals. All bouts occurred in social or feeding groups with the presence of calves (Table III). However, these sounds were generally rare. They were identified during only 10 of 82 *Inia* encounters, which comprised over 43 h of recordings across a wide variety of behavioral contexts and locations. Furthermore, downsweeps were not detected by Melo-Santos *et al.* (2019) in 15 h of recordings of a socializing/feeding group of human-habituated dolphins in the lower reaches of the Tocantins River. This suggests such sounds are used in specific behavioral contexts. Nevertheless, our GLMs did not show a statistically significant relationship between the emission of downsweeps and behavioral state and/or discrete behavioral events. The occurrence of bouts was correlated with long recording sessions, which might be related to the fact that foraging and socializing groups of *Inia* were

easier to track since they tended to stay in the same area for longer periods when engaged in these behaviors.

Downsweeps have been documented across several species of cetaceans: botos (*Inia geoffrensis*; May-Collado and Wartzok, 2007), Guiana dolphins *Sotalia guianensis* and tucuxis (*Sotalia fluviatilis*; Melo-Santos, 2018; Pivari and Rosso, 2005), pilot whales (*G. macrorhynchus* and *G. melas*; Dreher and Evans, 1964; Taruski, 1979), *Stenella longirostris* (Bazúa-Durán and Au, 2002), common dolphins (*Delphinus* species; Dreher and Evans, 1964; Ansmann *et al.*, 2007; Petrella *et al.*, 2012), bottlenose dolphin (*Tursiops truncatus*; Dreher and Evans, 1964; Janik *et al.*, 1994; Janik and Slater, 1998), orcas (*Orcinus orca*; Filatova *et al.*, 2012; Simonis *et al.*, 2012; Samarra *et al.*, 2015), humpback dolphins (*Sousa chinensis*; Van Parijs and Corkeron, 2001), franciscanas (*P. blainvillei*; Cremer *et al.*, 2017), and belugas (*Delphinapterus leucas*; Garland *et al.*, 2015). Among these studies Garland *et al.* (2015), May-Collado and Wartzok (2007), Petrella *et al.* (2012), and Samarra *et al.* (2015) reported downsweeps as the most common whistle type found in their samples. However, none of the above studies report downsweeps being used in a repetitive fashion or identify the contexts in which these signals were used. The exception is Dreher and Evans (1964), who reported that three juvenile bottlenose dolphins produced downsweeps in rapid repetition in situations of stress or fright. Our surface observations did not suggest that dolphins were in stress or a frightful situation (e.g., aggressive behavior) when downsweep bouts were produced. However, we cannot account for behavior of dolphins while submerged. Simonis *et al.* (2012) also noted the repetitive use of downsweep sounds by orcas, and because of their similarities to bat echolocation calls these authors




FIG. 4. (Color online) Visualization of downsweep characteristics in multivariate space according to two components in a PCA. Colours and shades indicate recording sessions. The “L” in group ID indicates animals recorded in the Lower Tocantins River, while the “M” stands for recordings of the Middle Tocantins River.

TABLE IV. Estimated parameters relating the presence of downsweep vocalizations in recordings of *Inia* to behavioral context and group characteristics. Relationships were estimated with a GLM (binomial family, logit-link).

	Estimate	SE	z	p
Intercept	-3.737	0.875	-4.270	<0.001
Behavior: Socializing	1.310	0.898	1.458	0.145
Behavior: Travelling	1.426	1.353	1.054	0.292
Number of calves	0.525	0.771	0.682	0.495
Group size	-0.149	0.111	-1.344	0.179
Recording duration	0.0412	0.018	2.330	0.020

TABLE V. Estimated parameters relating the presence of downsweep vocalizations in recordings of *Inia* to discrete behavioral events. Relationships were estimated with a GLM (binomial family, logit-link).

	Estimate	SE	z	p
Intercept	-3.746	0.817	-4.585	<0.001
Body contact	0.014	1.257	0.011	0.991
Flippers on belly	0.761	1.255	0.606	0.544
Leap	0.950	1.043	0.911	0.362
Object carrying	-0.249	1.799	-0.138	0.890
Penis exhibition	17.529	2399.545	0.007	0.994
Synchronized leap	-0.225	1.367	-0.165	0.869
Tail slap	-0.427	1.412	-0.302	0.763
Recording duration	0.025	0.014	1.801	0.072

suggested that orcas might use a series of downsweeps for echolocation. The downsweeps we recorded were much lower in frequency, and clicks were present continuously in all of our recordings, suggesting it is unlikely downsweeps serve as echolocation sounds. Moreover, if *Inia* downsweeps were used for echolocation one would expect them to be more common since echolocation is vital for *Inia* navigation and orientation. May-Collado and Wartzok (2007) suggested that botos use whistles primarily to maintain inter-individual distance. Our observations of botos producing downsweep bouts during social and foraging activities with synchronized surface behavior suggest that this might not be the case for Araguaian botos. Further support for such a functional difference comes from the fact that downsweeps presented in May-Collado and Wartzok (2007) for botos (*I. geoffrensis*) in the Napo River (Ecuador) appeared to have different frequency contours of those recorded in our study.

Cetaceans often use repeated sequences of sounds to broadcast their individual or group identity (Janik *et al.*, 2013; Gero *et al.*, 2016; Sayigh *et al.*, 2013). Identity is conveyed by the patterns of frequency modulation of sounds (Janik *et al.*, 2006) or through the order in which pulses are repeated (Watkins and Schevill, 1977). As boto downsweeps have relatively simple contours (Fig. 2) it is unlikely that they serve as individual identifiers in this way. However, two groups produced bouts of distinct downsweeps, suggesting possible group specificity. Nevertheless, their rare occurrence in our recordings makes it unlikely that they are required to maintain group cohesion as in delphinids (Janik, 2009). More common signals, such as pulsed calls (Melo-Santos *et al.*, 2019), likely play a more important role in social cohesion than downsweeps.

Given that downsweep emissions often occurred in long bouts (Fig. 3), and these bouts always occurred in the presence of calves, it is possible that they could function as begging calls (Godfray 1995a,b). Animal begging calls are normally produced by infants demanding resources (often food) from their parents (Godfray 1995a,b; Manser *et al.*, 2008). In cetaceans, sounds occurring before suckling have been recorded for humpback whales (*M. novaeangliae*); however, mechanical cues are more important to initiate nursing (Videsen *et al.*, 2017). Sounds associated with

suckling were also reported for captive neonate bottlenose dolphins (Morisaka *et al.*, 2005). In botos, sound bouts, especially whistles, might be important to start nursing. However, we did not detect a significant statistical relationship between the presence of calves and the production of downsweeps, possibly due to the small number of observations in our study.

In two out of five encounters with downsweep bouts, we also noticed surface displays which might indicate socio-sexual behavior such as object-carrying, exhibition of penis, and close contact between individuals. Mating calls of other animal species are also produced in bouts so as to attract the attention of possible mating partners, and this behavior is spread over a variety of taxa, including insects, amphibians, birds, and mammals (Catchpole and Slater, 2008; Fellers, 1979; Hedrick, 1986; Reby and Charlton, 2012). It is possible that Araguaian boto downsweeps have a similar function in the context of mating. Both downsweep production and social-sexual behavior are not observed very often, which makes this a difficult relationship to study. Future studies should aim to localize sounds to individuals to help with the identification of calling contexts.

ACKNOWLEDGMENTS

We are thankful to René Swift for the comments on the manuscript and designing the map. We thank Mark Johnson for lending us the SoundTrap used in this research. We thank Angélica Rodrigues, Danilo Arcoverde, Renata Saraiva, Alessandra Formento, Doracele Tuma, Adriano Tavares, and Jairo Azevedo for the help in fieldwork. Funding was provided by the Swarovski Foundation and World Wide Fund for Nature Brazil. We are also thankful to the community of the Capim Island for the support to our team in the field.

¹See <https://github.com/rflachlan/Luscinia/wiki> (Last viewed October 2019).

Altmann, J. (1974). "Observational study of behavior: Sampling methods," *Behaviour* **49**, 227–266.

Amorim, T. O. S., Andriolo, A., Reis, S. S., and Santos, M. E. (2016). "Vocalizations of Amazon river dolphins (*Inia geoffrensis*): Characterization, effect of physical environment and differences between populations," *J. Acoust. Soc. Am.* **139**, 1285–1293.

Ansmann, I. C., Goold, J. C., Evans, P. G., Simmonds, M., and Keith, S. G. (2007). "Variation in the whistle characteristics of short-beaked common dolphins, *Delphinus delphis*, at two locations around the British Isles," *J. Mar. Biol. Assoc. U. K.* **87**, 19–26.

Au, W. W., Popper, A. N., and Fay R. R., eds. (2000). *Hearing by Whales and Dolphins* (Springer, London, UK), pp. 1–485.

Bazúa-Durán, C., and Au, W. W. (2002). "The whistles of Hawaiian spinner dolphins," *J. Acoust. Soc. Am.* **112**, 3064–3072.

Best, R. C., and da Silva, V. M. F. (1989). "Amazon river dolphin, boto *Inia geoffrensis* (de Blainville, 1817)," in *Handbook of Marine Mammals*, edited by S. H. Ridgway, R. Harrison, and R. J. Harrison (Elsevier, London: Academic Press), Vol. 4, pp. 1–23.

Best, R. C., and da Silva, V. M. F. (1993). "*Inia geoffrensis*," *Mamm. Species* **426**, 1–8.

Buck, J. R., and Tyack, P. L. (1993). "A quantitative measure of similarity for *Tursiops truncatus* signature whistles," *J. Acoust. Soc. Am.* **94**, 2497–2506.

- Caldwell, M. C., Caldwell, D. K., and Evans, W. E. (1966). "Sounds and behavior of captive Amazon freshwater dolphins, *Inia geoffrensis*," Los Angeles Count. Mus. Nat. Hist. **108**, 1–24.
- Catchpole, C. K., and Slater, P. J. (2008). *Bird Song: Biological Themes and Variations*, 2nd ed. (Cambridge University Press, Cambridge, UK), pp. 1–355.
- Cremer, M. J., Holz, A. C., Bordino, P., Wells, R. S., and Simões-Lopes, P. C. (2017). "Social sounds produced by Franciscana dolphins, *Pontoporia blainvillei* (Cetartiodactyla, Pontoporiidae)," *J. Acoust. Soc. Am.* **141**, 2047–2054.
- Deecke, V. B., Barrett-Lennard, L. G., Spong, P., and Ford, J. K. (2010). "The structure of stereotyped calls reflects kinship and social affiliation in resident killer whales (*Orcinus orca*)," *Naturwissenschaften* **97**, 513–518.
- Deecke, V. B., and Janik, V. M. (2006). "Automated categorization of bioacoustic signals: Avoiding perceptual pitfalls," *J. Acoust. Soc. Am.* **119**, 645–653.
- Diazgranados, M. C., and Trujillo, F. (2002). "Vocal repertoire of the freshwater dolphins *Inia geoffrensis* and *Sotalia fluviatilis* in Colombia, South America," *J. Acoust. Soc. Am.* **112**, 2400.
- Ding, W., Würsig, B., and Evans, W. (1995). "Comparisons of whistles among seven odontocete species," in *Sensory Systems of Aquatic Mammals*, edited by R. A. Kastelein, J. A. Thomas, and P. E. Nachtigall (De Spil, Woerden, The Netherlands), pp. 299–323.
- Ding, W., Würsig, B., and Leatherwood, S. (2001). "Whistles of boto, *Inia geoffrensis*, and tucuxi, *Sotalia fluviatilis*," *J. Acoust. Soc. Am.* **109**, 407–411.
- Dreher, J. J., and Evans, W. E. (1964). "Cetacean communication," in *Marine Bioacoustics, 1*, edited by W. N. Tavolga (Oxford Pergamon, Symposium Publications Division), pp. 373–393.
- Duarte de Figueiredo, L., and Simão, S. M. (2009). "Possible occurrence of signature whistles in a population of *Sotalia guianensis* (Cetacea, Delphinidae) living in Sepetiba Bay, Brazil," *J. Acoust. Soc. Am.* **126**, 1563–1569.
- Fearey, J., Elwen, S. H., James, B. S., and Gridley, T. (2019). "Identification of potential signature whistles from free-ranging common dolphins (*Delphinus delphis*) in South Africa," *Anim. Cogn.* **22**, 777–789.
- Fellers, G. M. (1979). "Aggression, territoriality, and mating behaviour in North American treefrogs," *Anim. Behav.* **27**, 107–119.
- Filatova, O. A., Ford, J. K., Matkin, C. O., Barrett-Lennard, L. G., Burdin, A. M., and Hoyt, E. (2012). "Ultrasonic whistles of killer whales (*Orcinus orca*) recorded in the North Pacific (L)," *J. Acoust. Soc. Am.* **132**, 3618–3621.
- Ford, J. K. (1989). "Acoustic behaviour of resident killer whales (*Orcinus orca*) off Vancouver Island, British Columbia," *Can. J. Zool.* **67**, 727–745.
- Garland, E. C., Castellote, M., and Berchok, C. L. (2015). "Beluga whale (*Delphinapterus leucas*) vocalizations and call classification from the eastern Beaufort Sea population," *J. Acoust. Soc. Am.* **137**, 3054–3067.
- Gero, S., Whitehead, H., and Rendell, L. (2016). "Individual, unit and vocal clan level identity cues in sperm whale codas," *R. Soc. Open Sci.* **3**, 1–11.
- Godfray, H. C. J. (1995a). "Evolutionary theory of parent-offspring conflict," *Nature* **376**, 133–138.
- Godfray, H. C. J. (1995b). "Signaling of need between parents and young: Parent-offspring conflict and sibling rivalry," *Am. Nat.* **146**, 1–24.
- Gomez-Salazar, C., Trujillo, F., Portocarrero-Aya, M., and Whitehead, H. (2012a). "Population, density estimates, and conservation of river dolphins (*Inia* and *Sotalia*) in the Amazon and Orinoco river basins," *Mar. Mammal Sci.* **28**, 124–153.
- Gomez-Salazar, C., Trujillo, F., and Whitehead, H. (2012b). "Ecological factors influencing group sizes of river dolphins (*Inia geoffrensis* and *Sotalia fluviatilis*)," *Mar. Mammal Sci.* **28**, E124–E142.
- Goulding, M., Barthem, R., Ferreira, E. J. G., and Duenas, R. (2003). *The Smithsonian atlas of the Amazon* (Smithsonian Books, Washington), pp. 1–253.
- Hedrick, A. V. (1986). "Female preferences for male calling bout duration in a field cricket," *Behav. Ecol. Sociobiol.* **19**, 73–77.
- Hrbek, T., da Silva, V. M. F., Dutra, N., Gravena, W., Martin, A. R., and Farias, I. P. (2014). "A new species of river dolphin from Brazil or: How little do we know our biodiversity," *PLoS One* **9**, e83623.
- Janik, V. M. (2000). "Food-related bray calls in wild bottlenose dolphins (*Tursiops truncatus*)," *Proc. R. Soc. Lond. Ser. B* **267**, 923–927.
- Janik, V. M. (2005). "Underwater acoustic communication networks in marine mammals," in *Animal Communication Networks*, edited by P. K. McGregor (Cambridge University Press, Cambridge, UK), pp. 390–415.
- Janik, V. M. (2009). "Acoustic communication in delphinids," *Adv. Study Behav.* **40**, 123–157.
- Janik, V. M., King, S. L., Sayigh, L. S., and Wells, R. S. (2013). "Identifying signature whistles from recordings of groups of unrestrained bottlenose dolphins (*Tursiops truncatus*)," *Mar. Mammal Sci.* **29**, 109–122.
- Janik, V. M., Sayigh, L. S., and Wells, R. S. (2006). "Signature whistle shape conveys identity information to bottlenose dolphins," *Proc. Natl. Acad. Sci. U.S.A.* **103**, 8293–8297.
- Janik, V. M., and Slater, P. J. B. (1998). "Context-specific use suggests that bottlenose dolphin signature whistles are cohesion calls," *Anim. Behav.* **56**, 829–838.
- Janik, V. M., Todt, D., and Dehnhardt, G. (1994). "Signature whistle variations in a bottlenosed dolphin, *Tursiops truncatus*," *Behav. Ecol. Sociobiol.* **35**, 243–248.
- Jensen, F. H., Beedholm, K., Wahlberg, M., Bejder, L., and Madsen, P. T. (2012). "Estimated communication range and energetic cost of bottlenose dolphin whistles in a tropical habitat," *J. Acoust. Soc. Am.* **131**, 582–592.
- Junk, W. J., Piedade, M. T. F., Schöngart, J., Cohn-Haft, M., Adeney, J. M., and Wittmann, F. (2011). "A classification of major naturally-occurring Amazonian lowland wetlands," *Wetlands* **31**, 623–640.
- Kammaing, C., Van Hove, M. T., Englesma, F. J., and Terry, R. P. (1993). "Investigations on cetacean sonar X: A comparative analysis of underwater echolocation clicks of *Inia* spp. and *Sotalia* spp.," *Aquat. Mamm.* **19**, 31–43.
- Kaplan, M. B., Aran Mooney, T., Sayigh, L. S., and Baird, R. W. (2014). "Repeated call types in Hawaiian melon-headed whales (*Peponocephala electra*)," *J. Acoust. Soc. Am.* **136**, 1394–1401.
- Manser, M. B., Madden, J. R., Kunc, H. P., English, S., and Clutton-Brock, T. (2008). "Signals of need in a cooperatively breeding mammal with mobile offspring," *Anim. Behav.* **76**, 1805–1813.
- Marcoux, M., Auger-Méthé, M., and Humphries, M. M. (2012). "Variability and context specificity of narwhal (*Monodon monoceros*) whistles and pulsed calls," *Mar. Mammal Sci.* **28**, 649–665.
- Martin, A. R., Da Silva, V. M. F., and Rothery, P. (2008). "Object carrying as socio-sexual display in an aquatic mammal," *Biol. Lett.* **4**, 243–245.
- Martin, A. R., Silva, V. M. F., and Salmon, D. L. (2004). "Riverine habitat preferences of botos (*Inia geoffrensis*) and tucuxis (*Sotalia fluviatilis*) in the central Amazon," *Mar. Mammal Sci.* **20**, 189–200.
- May-Collado, L. J., Agnarsson, I., and Wartzok, D. (2007). "Phylogenetic review of tonal sound production in whales in relation to sociality," *BMC Evol. Biol.* **7**, 1–20.
- May-Collado, L. J., and Wartzok, D. (2007). "The freshwater dolphin *Inia geoffrensis geoffrensis* produces high frequency whistles," *J. Acoust. Soc. Am.* **121**, 1203–1212.
- McComb, K. E. (1991). "Female choice for high roaring rates in red deer, *Cervus elaphus*," *Anim. Behav.* **41**, 79–88.
- Melo-Santos, G. (2018). "Acoustic ecology of dolphins of the genus *Sotalia* (Cetartiodactyla, Delphinidae) and of the newly described Araguaian boto *Inia araguaiaensis* (Cetartiodactyla, Iniidae)," Ph.D. thesis, Universidade Federal do Pará, Belém, Pará, Brazil, available at <http://ppgtpc.proesp.ufpa.br/ARQUIVOS/dissertacoes/Gabriel%20Melo%20Santos.pdf> (Last viewed November, 2019).
- Melo-Santos, G., Rodrigues, A. L. F., Tardin, R. H., de Sá, M. I., Marmontel, M., Da Silva, M. L., and May-Collado, L. J. (2019). "The newly described Araguaian river dolphins, *Inia araguaiaensis* (Cetartiodactyla, Iniidae), produce a diverse repertoire of acoustic signals," *PeerJ* **7**, e6670.
- Mizue, K. A., Nishiwaki, M. A., and Takemura, A. K. (1971). "The underwater sound of Ganges river dolphins (*Platanista gangetica*)," *Sci. Rep. Whales Res. Inst.* **23**, 123–128.
- Morisaka, T., Shinohara, M., and Taki, M. (2005). "Underwater sounds produced by neonatal bottlenose dolphins (*Tursiops truncatus*): II. Potential function," *Aquat. Mamm.* **31**, 258–265.
- Overstrom, N. A. (1983). "Association between burst-pulse sounds and aggressive behavior in captive Atlantic bottlenosed dolphins (*Tursiops truncatus*)," *Zoo Biol.* **2**, 93–103.
- Pavanato, H. J., Melo-Santos, G., Lima, D. S., Portocarrero-Aya, M., Paschoalini, M., Mosquera, F., Trujillo, F., Meneses, R., Marmontel, M.,

- and Maretti, C. (2016). "Risks of dam construction for South American river dolphins: A case study of the Tapajós River," *Endanger. Species Res.* **31**, 47–60.
- Payne, R. S., and McVay, S. (1971). "Songs of humpback whales," *Science* **173**, 587–597.
- Penner, R. H., and Murchison, A. E. (1970). "Experimentally demonstrated echolocation in the Amazon River porpoise, *Inia geoffrensis* (Blainville)," in *Proc. 7th Ann. Conf. Biol. Sonar Diving Mammals*, Vol. 7, pp. 1–22.
- Pérez, J. M., Jensen, F. H., Rojano-Doñate, L., and Aguilar de Soto, N. (2017). "Different modes of acoustic communication in deep-diving short-finned pilot whales (*Globicephala macrorhynchus*)," *Mar. Mammal Sci.* **33**, 59–78.
- Petrella, V., Martínez, E., Anderson, M. G., and Stockin, K. A. (2012). "Whistle characteristics of common dolphins (*Delphinus* sp.) in the Hauraki Gulf, New Zealand," *Mar. Mammal Sci.* **28**, 479–496.
- Pivari, D., and Rosso, S. (2005). "Whistles of small groups of *Sotalia fluviatilis* during foraging behavior in southeastern Brazil," *J. Acoust. Soc. Am.* **118**, 2725–2731.
- Podos, J., Da Silva, V. M. F., and Rossi-Santos, M. R. (2002). "Vocalizations of Amazon river dolphins, *Inia geoffrensis*: Insights into the evolutionary origins of delphinid whistles," *Ethology* **108**, 601–612.
- R Core Team (2017). "R: A language and environment for statistical computing (version 3.4.2) [computer program]," Vienna: R Foundation for Statistical Computing, available at <https://www.Rproject.org/> (Last viewed July 2019).
- Rankin, S., Oswald, J., Barlow, J., and Lammers, M. (2007). "Patterned burst-pulse vocalizations of the northern right whale dolphin, *Lissodelphis borealis*," *J. Acoust. Soc. Am.* **121**, 1213–1218.
- Reby, D., and Charlton, B. D. (2012). "Attention grabbing in red deer sexual calls," *Anim. Cogn.* **15**, 265–270.
- Ribeiro, M. C. L. D. B., Petere, M., and Juras, A. A. (1995). "Ecological integrity and fisheries ecology of the Araguaia—Tocantins River Basin, Brazil," *River Res. Applic.* **11**, 325–350.
- Richardson, W. J., Greene, C. R., Jr., Malme, C. I., and Thomson, D. H. (1995). *Marine Mammals and Noise* (Academic, San Diego, CA), pp. 1–576.
- Sakoe, H., and Chiba, S. (1978). "Dynamic-programming algorithm optimization for spoken word recognition," *IEEE Trans. Acoust. Speech* **26**, 43–49.
- Samarra, F. I., Deecke, V. B., Simonis, A. E., and Miller, P. J. (2015). "Geographic variation in the time-frequency characteristics of high-frequency whistles produced by killer whales (*Orcinus orca*)," *Mar. Mammal Sci.* **31**, 688–706.
- Santos, G. M., and Jégu, M. (1989). "Inventário taxonômico e redescricao das espécies de anostomídeos (Characiformes, Anostomidae) do baixo rio Tocantins, PA, Brasil" ("Taxonomic inventory and redescription of anostomid (Characiformes, Anostomidae) species of the Lower Tocantins River"), *Acta Amazonica* **19**, 159–213.
- Santos, G. M. A., Quaresma, A. C., Barata, R. R., Martins, B. M., Siciliano, S., Silva, J. D. S., and Emin-Lima, R. (2012). "Etho-ecological study of the Amazon River dolphin, *Inia geoffrensis* (Cetacea: Iniidae), and the dolphins of the genus *Sotalia* (Cetacea: Delphinidae) in Guamá River, Amazonia," *Mar. Biodivers. Rec.* **5**, 1–5.
- Santos, G. M. A., Rodrigues, A. L. F., Arcoverde, D. L., Ramos, I., Sena, L., and Silva, M. L. (2014). "Unusual records of the behavior of boto *Inia* sp. (Cetartiodactyla, Iniidae) in the lower reaches of the Tocantins and Guamá Rivers, Amazônia," in *Dolphins: Ecology, Behavior and Conservation Strategies*, edited J. B. Samuels (Nova, New York), pp. 1–17.
- Sayigh, L., Quick, N., Hastie, G., and Tyack, P. L. (2013). "Repeated call types in short-finned pilot whales, *Globicephala macrorhynchus*," *Mar. Mammal Sci.* **29**, 312–324.
- Sayigh, L. S., Tyack, P. L., Wells, R. S., Solow, A. R., Scott, M. D., and Irvine, A. B. (1999). "Individual recognition in wild bottlenose dolphins: A field test using playback experiments," *Anim. Behav.* **57**, 41–50.
- Schevill, W. E., and Watkins, W. A. (1966). "Sound structure and directionality in *Orcinus* (killer whale)," *Zoologica* **51**, 71–76.
- Simonis, A. E., Baumann-Pickering, S., Oleson, E., Melcón, M. L., Gassmann, M., Wiggins, S. M., and Hildebrand, J. A. (2012). "High-frequency modulated signals of killer whales (*Orcinus orca*) in the North Pacific," *J. Acoust. Soc. Am.* **131**, 295–301.
- Slater, P. J. B., and Lester, N. P. (1982). "Minimising errors in splitting behaviour into bouts," *Behaviour* **79**, 153–161.
- Smith, J. N., Goldizen, A. W., Dunlop, R. A., and Noad, M. J. (2008). "Songs of male humpback whales, *Megaptera novaeangliae*, are involved in intersexual interactions," *Anim. Behav.* **76**, 467–477.
- Smolker, R. A., Mann, J., and Smuts, B. B. (1993). "Use of signature whistles during separations and reunions by wild bottlenose dolphin mothers and infants," *Behav. Ecol. Sociobiol.* **33**, 393–402.
- Taruski, A. G. (1979). "The whistle repertoire of the North Atlantic pilot whale (*Globicephala melaena*) and its relationship to behavior and environment," in *Behavior of Marine Animals*, edited by H. E. Winn and B. L. Olla (Springer, Boston, MA), pp. 345–368.
- Tyack, P. L. (2000). "Functional aspects of cetacean communication," in *Cetacean Societies: Field Studies of Dolphins and Whales*, edited by J. Mann, R. C. Connor, P. L. Tyack, and H. Whitehead (The University of Chicago Press, Chicago, IL), pp. 270–307.
- Van Parijs, S. M., and Corkeron, P. J. (2001). "Vocalizations and behaviour of Pacific humpback dolphins *Sousa chinensis*," *Ethology* **107**, 701–716.
- Videsen, S. K., Bejder, L., Johnson, M., and Madsen, P. T. (2017). "High suckling rates and acoustic crypsis of humpback whale neonates maximize potential for mother-calf energy transfer," *Funct. Ecol.* **31**, 1561–1573.
- Wang, D., Wang, K. X., and Akamatsu, T. K. (1999). "Study on whistling of the Chinese river dolphin (*Lipotes vexillifer*)," *Oceanol. Limnol. Sin.* **30**, 353–360.
- Wang, K., Wang, D., Akamatsu, T., Fujita, K., and Shiraki, R. (2006). "Estimated detection distance of a baiji's (Chinese river dolphin, *Lipotes vexillifer*) whistles using a passive acoustic survey method," *J. Acoust. Soc. Am.* **120**, 1361–1365.
- Wang, D., Wenxiang, L., and Zhifan, W. (1995). "A preliminary study of the acoustic behavior of the baiji, *Lipotes vexillifer*," in *Biology and Conservation of the River Dolphins* (IUCN Special Survival Commission, Gland, Switzerland), pp. 137–140.
- Watkins, W. A., and Schevill, W. E. (1977). "Sperm whale codas," *J. Acoust. Soc. Am.* **62**, 1485–1490.
- Xiao, Y., and Jing, R. (1989). "Underwater acoustic signals of the baiji, *Lipotes vexillifer*," in *Biology and Conservation of the River Dolphins, Occasional paper* (IUCN Special Survival Commission, Gland, Switzerland), Vol. 3, pp. 137–140.
- Zwamborn, E. M. J., and Whitehead, H. (2017). "Repeated call sequences and behavioural context in long-finned pilot whales off Cape Breton, Nova Scotia, Canada," *Bioacoustics* **26**, 169–183.